

Research Article

Salinity tolerance of non-native suckermouth armoured catfish (Loricariidae: *Pterygoplichthys* sp.) from Kerala, India

A. Biju Kumar¹, Pamela J. Schofield^{2,*}, Smrithy Raj¹ and Sima Satheesh¹

¹Department of Aquatic Biology & Fisheries, University of Kerala, Thiruvananthapuram 695581, India

²US Geological Survey, Wetland and Aquatic Research Center, Gainesville, FL 32653, USA

*Corresponding author

E-mail: pschofield@usgs.gov

Received: 16 June 2017 / Accepted: 15 November 2017 / Published online: 4 December 2017

Handling editor: Vadim E. Panov

Abstract

Loricariid catfishes of the genus *Pterygoplichthys* are native to South America and have been introduced in many localities around the world. They are freshwater fishes, but may also use low-salinity habitats such as estuaries for feeding or dispersal. Here we report results of a field survey and salinity-tolerance experiments for a population of *Pterygoplichthys* sp. collected in Kerala, India. In both chronic and acute salinity-tolerance trials, fish were able to withstand salinities up to 12 ppt with no mortality; however, fish transferred to salinities > 12 ppt did not survive. The experimental results provide evidence that non-native *Pterygoplichthys* sp. are able to tolerate mesohaline conditions for extended periods, and can easily invade the brackish water ecosystems of the state. Further, *Pterygoplichthys* sp. from Kerala have greater salinity tolerance than other congeners. These data are vital to predicting the invasion of non-native fishes such as *Pterygoplichthys* spp. into coastal systems in Kerala and worldwide. This is particularly important as estuarine ecosystems are under threat of global climate change and sea-level rise. In light of the results of the present study and considering the reports of negative impacts of the species in invaded water bodies, management authorities may consider controlling populations and/or instituting awareness programmes to prevent the spread of this nuisance aquatic invasive species in Kerala.

Key words: ecophysiology, estuary, invasion, non-native fish, osmoregulation

Introduction

Trade in aquarium fishes is one of the major pathways for the invasion of potentially harmful aquatic invasions globally (Gozlan et al. 2010; Magalhães and Jacobi 2013; Emiroğlu et al. 2016; Allen et al. 2017; Marr et al. 2017). The ever-increasing trade of ornamental fishes is one of the most important pathways for aquatic invasion in India (Singh et al. 2013; Raghavan et al. 2013; Kumar et al. 2015; Soundararajan et al. 2015; Sandilyan 2016; Muralidharan 2017). Among commonly-traded freshwater ornamental fishes, suckermouth armoured catfishes of the family Loricariidae and genus *Pterygoplichthys* T.N. Gill native to inland water bodies of South America, have established successful invasive populations throughout the world (Fuller et al. 1999; Nico and Martin 2001; Nakabo

2002; Chavez et al. 2006; Joshi 2006; Page and Robins 2006; Armando et al. 2007; Ozdilek 2007; Hossain et al. 2008; Keszka et al. 2008; Nico et al. 2009; Piazzini et al. 2010; Simonović et al. 2010; Wu et al. 2011; Gibbs et al. 2013; Golani and Snovsky 2013; Jones et al. 2013; Rueda-Jasso et al. 2013; Sumanasinghe and Amarasinghe 2013; Zworykin and Budaev 2013; Ishikawa and Tachihara 2014; Jumawan and Herrera 2014; Wei et al. 2017). *Pterygoplichthys* spp. are sold in markets under various common names including armoured catfishes, janitor fish, sailfin catfishes, plecos, glass cleaner catfishes, suckermouth catfishes and others. In India *Pterygoplichthys* spp. have been recorded from several natural water bodies and in many places they have become invasive species within the last two decades (Daniels 2006; Krishnakumar et al. 2009; Knight 2010; Sinha et al.

2010; Singh 2014; Singh et al. 2013, 2014; Kumar et al. 2015; Muralidharan et al. 2015; Panikkar et al. 2015; Meena et al. 2016; Rao and Sunchu 2017).

Pterygoplichthys catfishes have many attributes that are thought to have facilitated their invasion around the world. They are able to tolerate wide fluctuations in flow regimes (Welcomme and Vidthayanom 2003; Nico et al. 2012) and resist hypoxia due to structural modifications in the stomach that allow them to breathe atmospheric air (da Cruz et al. 2013). They also have efficient reproductive strategies and deter predation by possessing bony plates that cover their body (Hoover et al. 2004; Wei et al. 2017). Furthermore, invasion success of a non-native species is largely determined by its ability to adapt to the physiochemical parameters in the invaded environment (Kestrup and Ricciardi 2010). For many non-native freshwater fishes, salinity is an important factor that prevents the invasion or acts as a barrier against dispersal (Scott et al. 2008; Gutierrez et al. 2014). Salinity may also delimit an invasive species' pattern of dispersal, population structure, abundance and interaction with native species (Cognetti and Maltagliati 2000; Smith et al. 2004; Paavola et al. 2005; Alcaraz and Garcia-Berthou 2007; Alcaraz et al. 2008; Schofield and Nico 2009).

We were first made aware of the presence of non-native suckermouth sailfin catfishes of the genus *Pterygoplichthys* in Kerala state of India by local fishers and ornamental fish traders who collect juveniles for trade. The morphological identification of some of the commonly-traded species in the tropics such as *P. anisitsi*, *P. multiradiatus*, *P. pardalis* and *P. disjunctivus* are separated primarily based on the nature of abdominal patterns (Nico et al. 2012). However, the abdominal patterns may overlap in the case of hybrids, making the process of identification difficult or impossible. The species that has invaded Kerala appears to be a hybrid (Kumar et al. 2015).

The invasive populations of *Pterygoplichthys* spp. in many parts of the world threaten indigenous species and in many cases surpass the biomass of native species (Nico et al. 2009; Capps and Flecker 2013). In this study, we determine the salinity tolerance of *Pterygoplichthys* sp., which may help predict the ultimate range in Kerala. Herein, we provide details of a field effort to better survey the geographic range of these fishes in Kerala as well as experimental results from salinity-tolerance tests. Our results are compared to the few previous studies of salinity tolerance in *Pterygoplichthys* spp. and results from field collections in low-salinity waters.

Methods

Field survey and fish collection

The water bodies of Thiruvananthapuram district of Kerala state, India were surveyed during 2015–2016 to detect the presence of non-native *Pterygoplichthys* sp. Fortnightly collections were made from each water body using cast nets and gill nets. Surveys to collect the fishes were done in the fresh- and brackish-water inland ecosystems of southern Kerala including rivers (Karamana and Neyyar), drainage canals (Amayizhanchan Thodu), Vellayani freshwater lake and Veli brackish-water lake (Figure 1). Though both juveniles and adults were collected during the survey, only adult fishes were used for the salinity-tolerance experiments.

Acute salinity tolerance

Acute (abrupt) salinity tolerance experiments with *Pterygoplichthys* sp. were done following the protocols described by Kefford et al. (2004), Bringolf et al. (2005), Schofield and Nico (2009) and Capps et al. (2011), with some modifications. Fishes were collected from the natural drainage, Amayizhanchan Thodu (0 ppt salinity), Thiruvananthapuram city, Kerala. In total, 200 acclimatized *Pterygoplichthys* were selected for testing.

The experimental setup consisted of 20 (10-L) glass aquaria (30 cm long × 23 cm wide × 15 cm high) filled with 7 L of water. The tanks were allocated to ten different salinity treatments (0 [control], 2, 4, 8, 12, 16, 20, 24, 28 and 35 ppt; n = 2 replicates per treatment). Tanks were randomly arranged to avoid experimental artefacts. The salinity concentrations were prepared using both natural seawater and artificial sea salt (Red Sea® aquarium salt, Israel). After acclimatization, fishes were weighed (+/- 0.1 cm standard length [SL]), measured (+/- 1 g) and then randomly placed in the tanks (one fish per tank) to begin the experiment. Fish ranged in size from 181 to 229 mm SL (mean 208 mm) and 120 to 245 g mass (mean 182 g). Mass did not differ amongst treatments (one-way ANOVA, F[9,190] = 1.496; P = 0.15; Levene's test P = 0.89).

The fish were kept in the experimental tank for a period of 240 hours (10 days) or until death. All the tanks were covered with tight nylon meshes to prevent the fish from jumping out. As the experimental fish were air breathers, separate aeration was not attempted in the experimental tanks. They were not provided food during the experiment, as food could have fouled the water and this hardy fish can survive without food for at least a month (ABK pers. obs.). Survival was recorded every 15 minutes for the




Figure 1. Map of sites sampled for *Pterygoplichthys* in Kerala state, India. For details see Supplementary information Table S1.

first hour, every 1 hour for the first 24 hours, every 3 hours for the first 24–48 hours, and every 6 hours for the remaining period of experiment. Water temperature and salinity of each tank was monitored using a thermometer (± 0.1 °C) and salinometer (± 1 ppt). Water temperatures of all tanks were recorded each day throughout the experimental period. Temperatures ranged from 24.8 to 32.0 °C, but did not differ among salinities. The entire experiment was replicated 10 times, providing a total of 20 replicates for each treatment when data were combined (total = 200 fish).

Chronic salinity tolerance

To determine the species' tolerance to gradually-increased salinities, a chronic salinity tolerance experiment was completed. Thirty-two fish were assigned to one of eight treatments (0 [control], 2, 4, 6, 8, 10, 12 or 14 ppt); with four replicates tanks per salinity treatment. Tanks were randomly arranged to avoid experimental artefacts. Fish ranged in size from 76 to 245 mm SL (mean 191 mm) and 115 to 264 g mass (mean 1582 g). Mass did not differ amongst treatments (one-way ANOVA, $F[7,24] = 2.3434$; $P = 0.06$;

Table 1. Collection details for *Pterygoplichthys* sp. in various water bodies of Thiruvananthapuram district, Kerala.

Site	Habitat type	Salinity (ppt)	N	Length range (mm SL)
Neyyar River	Freshwater	0	0	
Karamana River	Freshwater	0	4	154–173
Vellayani lake	Freshwater	0	11	165–211
Veli lake	Brackish-water	0–8	3	188–194
Amayizhanchan Thodu	Freshwater	0	314	181–229
Total			332	




Figure 2. Acute salinity tolerance of *Pterygoplichthys* sp. from Kerala, India. Salinities tested include: 0 (control), 2, 4, 8, 12, 16, 20, 24, 28 and 35 ppt. Twenty replicates were tested for each salinity; total number of fish tested = 200. Fish were transferred directly from 0 ppt to test salinities and held there for 10 days (240 hours).

Levene's test $P < 0.001$). Fishes were held in 100-L aquaria and all started at 0 ppt. Salinities were then gradually increased (a change of 2 ppt every 2 days) for all treatments except the control until the treatment attained its target salinity. Survival was measured daily and dead fish, if any, were removed. At the end of the experiment, fish had been exposed to the control salinity for 23 days, 2 ppt for 20 days, 4 ppt for 17 days, 6 ppt for 14 days, 8 ppt for 11 days, 10 ppt for 8 days and 12 ppt for 5 days.

Data analysis

For both acute and chronic experiments, fish mass was compared across treatments with one-way Analysis of Variance (ANOVA) to ensure fish size did not differ amongst treatments. Levene's test was used to check for heteroscedasticity. Life expectancy was estimated with the Kaplan-Meier product-limit estimator (Kaplan and Meier 1958) and the log-rank test was used to compare survivorship curves (Savage 1956; Cox and Oakes 1984). For the acute-salinity challenge, all treatments began at the same time (time = 0). However, for the chronic-salinity experiment,

fish reached their target salinities sequentially (i.e., staggered over time). Thus, for the chronic-salinity experiment the day the fish reached their target salinity was designated as time = 0 for that treatment. We set our alpha level for statistical significance at 0.05. All data were analysed using NCSS version 11.

Results

Distribution of Pterygoplichthys sp.

The field surveys detected the presence of non-native *Pterygoplichthys* spp. from Karamana River, Amayizhanchan Thodu, Veli Lake and Vellayani Lake in the Thiruvananthapuram district of Kerala, India. We captured 332 individuals of *Pterygoplichthys* sp. from the water bodies of Thiruvananthapuram district, indicating the spread of species into various water bodies (Table 1). The survey results showed that Amayizhanchan Thodu, the natural drainage channels of Thiruvananthapuram District of Kerala, is heavily invaded by *Pterygoplichthys* sp., and the presence of this species in Veli Lake shows the ability of species to invade brackish waters of the state.

Acute salinity tolerance

No fish died in treatments ranging from 0–12 ppt during the 240 hrs (10 days) of the experiment. However, above 12 ppt survival decreased significantly (Figure 2, Table 2). Individuals lived about three hours at 16 ppt, two hours at 20 ppt and one hour at 24 ppt. At salinities of 28 and 35, fish died within one hour. Log-rank tests showed that each treatment of 16–35 ppt were distinctly different from all other treatments (Table 2).

Chronic salinity tolerance

Pterygoplichthys sp. survived all treatments from 0 to 12 ppt with no mortality. After reaching 14 ppt, individuals died in 24–36 hrs (mean = 30 hrs). It was not possible to compare treatments with the log-rank test as there was only mortality in one treatment (14 ppt).

Discussion

Many native and introduced freshwater fish species occasionally use estuarine and brackish water areas for feeding or dispersal (Loftus 1988; McIvor and Odum 1988; Brown et al. 2007). Fishes are limited to suitable salinities because of the stress exerted on osmoregulation by the surrounding water medium. This becomes an important aspect of biogeographic distribution (Myers 1949). Loricariids are a family of freshwater fishes (Nelson 1994) but have invaded estuarine environments in several countries beyond their native range due to their ability to tolerate low levels of salinity, as shown in this report.

Pterygoplichthys spp. possess several characteristics that enable them to establish invasive populations in their non-native range (Douglas et al. 2002; Ebenstein et al. 2015; Geerinckx et al. 2011; German and Bittong 2009; Harter et al. 2014; Jumawan and Herrera 2015; Villalba-Villalba et al. 2017; Zworykin and Budaev 2013). Some of these include: 1) the ability to survive anoxic conditions by means of accessory respiration with diverticula of the gastrointestinal tract (Armbruster 1998); 2) the presence of bony armor plates and spines, which makes them relatively difficult to consume; 3) reproductive advantages that facilitate invasion such as prolonged reproductive period, batch spawning, development of eggs at very low water levels and active parental care (Hoover et al. 2004; Liang et al. 2005); and 4) the ability to exploit ecologically-degraded aquatic habitats, which favor invasive species such as *Pterygoplichthys* spp. that can withstand poor water quality conditions (da Cruz et al. 2013; Parente et al. 2010).

Table 2. Results from acute salinity exposure experiments for *Pterygoplichthys* sp. A: Mean survival times (h) for each salinity and standard deviation (SD). * = unable to compute mean or SD as all fish survived the treatment for the entire length of the experiment (240 hrs). B: Log-rank tests comparing survival between treatments. Statistic given is *P*. Only treatments with mortality are compared (16–35 ppt).

A					
Salinity	Mean survival time (h)		SD		
0	*				
2	*				
4	*				
8	*				
12	*				
16	3.2		0.7		
20	2.3		0.47		
24	1.11		0.39		
28	0.5		0		
35	0.5		0		

B					
Salinity	16	20	24	28	35
16	*	< 0.001	< 0.001	< 0.001	< 0.001
20		*	< 0.001	< 0.001	< 0.001
24			*	< 0.001	< 0.001
28				*	NS
35					*

We have shown that *Pterygoplichthys* sp. is able to tolerate low levels of salinity, and that is yet another factor potentially explaining the invasion success of this group. In India, dumping of ornamental fishes is the primary route for introduction of non-native species into aquatic ecosystems (Kumar 2000; Singh and Lakra 2011; Soundararajan et al. 2015; Sandilyan 2016). The armoured catfishes of the genus *Pterygoplichthys* have been reported from various natural water bodies of India in the last decade, and in later stages as invasive populations in many major rivers and lakes (Knight 2010; Sinha et al. 2010; Singh and Lakra 2011; Singh et al. 2014; Kumar et al. 2015; Panikkar et al. 2015; Meena et al. 2016; Rao and Sunchu 2017). According to Kumar et al. (2015) the ornamental fish trade could be the reason for the introduction of *Pterygoplichthys* sp. in the water bodies of Kerala. This study shows that these catfishes have the potential to spread to other freshwater and brackish water ecosystems, possibly using brackish water “salt bridges” to reach new freshwater drainages.

Two studies have previously evaluated the salinity tolerance of *Pterygoplichthys* sp. using an experimental approach. In Mexico, Capps et al. (2011) recorded collections of *Pterygoplichthys* sp. from the field at salinities up to 8 ppt. The acute salinity tolerance of field-collected samples was tested, and it was found

that the species was able to survive abrupt (acute) exposure to salinities up to 9 ppt with little mortality over 10 days. Interestingly, the *Pterygoplichthys* sp. tested in Mexico appear slightly more sensitive to salinity than the ones in this study. Mexican *Pterygoplichthys* sp. survived an average of 58 hrs at 11 ppt and all fish at 12 ppt were dead by 40 hrs, whereas in our study all fish in salinities up to 12 ppt survived with no mortality for the entire length of the experiment (240 hours). Brion et al. (2013) reported non-native *Pterygoplichthys* sp. collected from the Philippines at salinities of 5–8 ppt. They conducted acute-salinity-tolerance tests on specimens purchased from a pet store, and found the LC₅₀ was 10.6 ppt for a 96-hour period. Interestingly, fish in salinity treatments of 15 and 20 ppt survived for an average of 47 and 24 hours, respectively. That was longer than the fish in the present study survived at these higher salinities. Because of difficulties in identification of *Pterygoplichthys* species, it is unclear whether the work of Brion et al. (2013) and Capps et al. (2011) was on the same species we have studied in Kerala. It is possible that different species have different tolerances to salinities; however, without better species discrimination it is not possible to address that question at this time. A few additional studies have recorded collections of non-native *Pterygoplichthys* sp. in low-salinity waters: Chavez et al. (2006) collected the fish in waters to 2 ppt in the Philippines. Golani and Snovsky (2013) collected *Pterygoplichthys pardalis* and *P. disjunctivus* in Israel at salinities from 0.5–2.0 ppt. Arévalo-Frías and Mendoza-Carranza (2015) reported *Pterygoplichthys* sp. in Mexico from 1.6–5.6 ppt. Finally, Idelberger et al. (2011) reported *Pterygoplichthys* spp. from Charlotte Harbor estuary (Florida, USA) at salinities up to at least 5 ppt and possibly as high as 12 ppt.

While *Pterygoplichthys* sp. are tolerant of low levels of salinity (to 12 ppt), higher salinities could limit migration through estuarine waters and confine the fish to its current distribution. Salinity may also limit further dispersal of this fish to other freshwater drainages through the marine route, although during monsoon flooding, when the estuaries, mangrove wetlands and rivers of India are flooded with freshwater, temporary aquatic connections may emerge (Selvam 2003; Martin et al. 2008). The major hydrological variable in the backwaters of Kerala is salinity, with a gradual decline in salinity from 30 ppt at the entrance of the estuary to 0.2 ppt at the confluence of the tributary and the rivers (Menon et al. 2000). Upstream of the river entry are farms built below-sea-level (Koohafkan and Altieri 2011), where traditional rice-fish culture and prawn filtration are have long been practiced (Subramanian 1988; Kurup and

Ranjeet 2002). In the estuaries and backwaters of Kerala during both southwest and northeast monsoons there is considerable flooding with freshwater, rapid decline in salinities across the habitat, and low salinities or even freshwater conditions in river mouths. These conditions may facilitate *Pterygoplichthys* sp. survival and dispersal in these regions.

The experimental results provided here show that non-native *Pterygoplichthys* sp. in Kerala are able to tolerate salinities up to 12 ppt for extended periods, and can invade brackish-water ecosystems of the state. The salinity tolerance of Kerala *Pterygoplichthys* sp. is greater than many typical freshwater fishes that can tolerate low levels of salinity (to about 8 ppt; Peterson and Meador 1994). These data are vital to predicting the invasion success and potential geographic spread of non-native fishes such as *Pterygoplichthys* spp. into coastal systems in Kerala and worldwide. Knowing salinity tolerance levels and predicting invasions are particularly important because invasive *Pterygoplichthys* catfishes have been reported to have negative effects on native fauna and local fisheries (Hoover et al. 2004; Sumanasinghe and Amarasinghe 2013) and management authorities may be able to control populations (e.g., Nico et al. 2012; Gibbs et al. 2013; Hill and Sowards 2015) or provide awareness programmes to prevent the spread of this nuisance species in both freshwater and estuarine habitats.

Acknowledgements

We thank the support of Directorate of Environment and Climate Change, Government of Kerala for the financial support to study aquatic invasives in Kerala state. Smrithy Raj thanks the government for the Rajiv Gandhi National Fellowship offered for the study. Rules for prevention of cruelty to animals in India exclude fishes, and the species used in the experiments is an invasive species that is culled to conserve the native fauna. We also thank the assistance of fisherman Mr Madhu for collection of specimens. K Reaver and D Gregoire-Lucente helped conduct literature reviews and format the report. Any use of trade, product or firm names is for descriptive purposes only and does not imply endorsement by the US Government.

References

- Alcaraz C, García-Berthou E (2007) Life history variation of an invasive mosquitofish (*Gambusia holbrooki*) along a salinity gradient. *Biological Conservation* 139: 83–92, <https://doi.org/10.1016/j.biocon.2007.06.006>
- Alcaraz C, Bisazza A, García-Berthou E (2008) Salinity mediates the competitive interactions between invasive mosquitofish and an endangered fish. *Oecologia* 155: 205–213, <https://doi.org/10.1007/s00442-007-0899-4>
- Allen PE, Barquero MD, Bermúdez E, Calderón JC, Hilje B, Pineda W, Saborio-Rodríguez G, Arguedas V, Chacón-Madriral E (2017) Calling for more accurate information in aquarium trade: analysis of live-fish import permits in Costa Rica. *Management of Biological Invasions* 8: 533–542, <https://doi.org/10.3391/mbi.2017.8.4.08>

- Arévalo-Frías W, Mendoza-Carranza M (2015) Influence of temporal and spatial factors on abundance and richness of fish early stages in shallow tropical estuaries. *Environmental Biology of Fishes* 98: 891–904, <https://doi.org/10.1007/s10641-014-0324-x>
- Armando TWK, Ramon RC, Enrique AA (2007) Amazon sailfin catfish *Pterygoplichthys pardalis* (Castelnau, 1885) (Loricariidae), another exotic species established in south-eastern Mexico. *Southwestern Naturalist* 52: 141–144, [https://doi.org/10.1894/0038-4909\(2007\)52\[141:ASCPPC\]2.0.CO;2](https://doi.org/10.1894/0038-4909(2007)52[141:ASCPPC]2.0.CO;2)
- Armbruster JW (1998) Modifications of the digestive tract for holding air in loricariid and scoloplacid catfishes. *Copeia* 3: 663–675, <https://doi.org/10.2307/1447796>
- Bringolf RB, Kwak TJ, Cope WG, Larimore MS (2005) Salinity tolerance of flathead catfish: implications for dispersal of introduced populations. *Transactions of the American Fisheries Society* 134: 927–936, <https://doi.org/10.1577/T04-195.1>
- Brion MA, Guillermo Jr JG, Uy C, Chavez J, Carandang IV JS (2013) Salinity tolerance of introduced South American sailfin catfishes (Loricariidae: *Pterygoplichthys* GILL 1858). *Philippine Journal of Science* 142(1): 13–19
- Brown JA, Scott DM, Wilson RW (2007) Do estuaries act as saline bridges to allow invasion of new freshwater systems by non-indigenous fish species? In: Gherardi F (ed), *Biological Invaders in Inland Waters: Profiles, Distribution and Threats*. Springer, Dordrecht, pp 401–414, https://doi.org/10.1007/978-1-4020-6029-8_21
- Capps KA, Flecker AS (2013) Invasive aquarium fish transform ecosystem nutrient dynamics. *Proceedings of the Royal Society B* 280: 20131520, <https://doi.org/10.1098/rspb.2013.1520>
- Capps KA, Nico LG, Mendoza-Carranza M, Arévalo-Frías W, Ropicki AJ, Heilpern SA, Rodiles-Hernández R (2011) Salinity tolerance of non-native suckermouth armoured catfish (Loricariidae: *Pterygoplichthys*) in south-eastern Mexico: implications for invasion and dispersal. *Aquatic Conservation: Marine and Freshwater Ecosystems* 21: 528–540, <https://doi.org/10.1002/aqc.1210>
- Chavez HM, Casao EA, Villanueva EP, Paras MP, Guinto MC, Mosqueda MB (2006) Heavy metal and microbial analyses of janitor fish (*Pterygoplichthys* spp.) in Laguna de Bay, Philippines. *Journal of Environmental Science and Management* 9: 31–40
- Cognetti G, Maltagliati F (2000) Biodiversity and adaptive mechanisms in brackish water fauna. *Marine Pollution Bulletin* 40: 7–14, [https://doi.org/10.1016/S0025-326X\(99\)00173-3](https://doi.org/10.1016/S0025-326X(99)00173-3)
- Cox DR, Oakes D (1984) *Analysis of Survival Data*. Monographs on Statistics and Applied Probability. Chapman and Hall, New York, 212 pp
- da Cruz AL, da Silva HR, Lundstedt LM, Schwantes AR, Moraes G, Klein W, Fernandes MN (2013) Air-breathing behavior and physiological responses to hypoxia and air exposure in the air-breathing loricariid fish, *Pterygoplichthys anisitsi*. *Fish Physiology and Biochemistry* 39: 243–256, <https://doi.org/10.1007/s10695-012-9695-0>
- Daniels RJR (2006) Introduced fishes: a potential threat to the native freshwater fishes of Peninsular India. *Journal of the Bombay Natural History Society* 103: 346–348
- Douglas RH, Collin SP, Corrigan J (2002) The eyes of suckermouth armoured catfish (Loricariidae, subfamily Hypostomus): pupil response, lenticular longitudinal spherical aberration and retinal topography. *Journal of Experimental Biology* 205: 3425–3433
- Ebenstein D, Calderon C, Troncoso OP, Torres FG (2015) Characterization of dermal plates from armoured catfish *Pterygoplichthys pardalis* reveals sandwich-like nano composite structure. *Journal of Mechanical Behavior of Biomedical Materials* 45: 175–182, <https://doi.org/10.1016/j.jmbm.2015.02.002>
- Emiroğlu O, Ekmekci FG, Aksu S, Baskurt S, Atalay A, Tarkan AS (2016) Introduction and establishment of tropical ornamental fish, *Pterygoplichthys* spp. (Actinopterygii: Siluriformes: Loricariidae) in hot springs: aquarium trade as a potential risk for biodiversity in Turkey. *Acta Ichthyologica et Piscatoria* 46: 351–356, <https://doi.org/10.3750/AIP2016.46.4.07>
- Fuller PL, Nico LG, Williams GD (1999) Nonindigenous fishes introduced into inland waters of the United States. *American Fisheries Society*, Special Publication 27: 435–441
- Geerinckx T, Herrel A, Adriaens D (2011) Suckermouth armoured catfish resolve the paradox of simultaneous respiration and suction attachment: a kinematic study of *Pterygoplichthys disjunctivus*. *Journal of Experimental Zoology. Part A, Ecological Genetics and Physiology* 315A: 121131, <https://doi.org/10.1002/jez.656>
- German DP, Bittong RA (2009) Digestive enzyme activities and gastrointestinal fermentation in wood-eating catfishes. *Journal of Comparative Physiology B: Biochemical, Systemic, and Environmental Physiology* 179: 1025–1042, <https://doi.org/10.1007/s00360-009-0383-z>
- Gibbs MA, Kurth BN, Bridges CD (2013) Age and growth of the loricariid catfish *Pterygoplichthys disjunctivus* in Volusia Blue Spring, Florida. *Aquatic Invasions* 8: 207–218, <https://doi.org/10.3391/ai.2013.8.2.08>
- Golani D, Snovsky G (2013) Occurrence of suckermouth armoured catfish (Siluriformes, Loricariidae, *Pterygoplichthys*) in inland waters of Israel. *BiolInvasion Records* 2: 253–256, <https://doi.org/10.3391/bir.2013.2.3.13>
- Gozlan RE, Britton JR, Cowx IG, Copp GH (2010) Current knowledge on non-native freshwater fish introductions. *Journal of Fish Biology* 76: 751–786, <https://doi.org/10.1111/j.1095-8649.2010.02566.x>
- Gutierrez SMM, Vitule JRS, Freire CA, Prodocimo V (2014) Physiological tools to predict invasiveness and spread via estuarine bridges: tolerance of Brazilian native and worldwide introduced freshwater fishes to increased salinity. *Marine and Freshwater Research* 65: 425–436, <https://doi.org/10.1071/MF13161>
- Harter TS, Shartau RB, Baker DW, Jackson DC, Val AL, Brauner CJ (2014) Preferential intracellular pH regulation represents a general pattern of pH homeostasis during acid-base disturbances in the armoured catfish, *Pterygoplichthys pardalis*. *Journal of Comparative Physiology B* 184: 709–718, <https://doi.org/10.1007/s00360-014-0838-8>
- Hill JE, Sowards J (2015) Successful eradication of the non-native loricariid catfish *Pterygoplichthys disjunctivus* from the Rainbow River, Florida. *Management of Biological Invasions* 6: 311–317, <https://doi.org/10.3391/mbi.2015.6.3.11>
- Hoover JJ, Killgore KJ, Cofrancesco AF (2004) Suckermouth catfishes: threats to aquatic ecosystems of the United States? *Aquatic Nuisance Species Research Program Bulletin* 4: 1–8
- Hossain MY, Rahman MM, Ahmed ZF, Ohtomi J, Islam ABMS (2008) First record of the South American sailfin catfish *Pterygoplichthys multiradiatus* in Bangladesh. *Journal of Applied Ichthyology* 24: 718–720
- Idelberger CF, Stafford CH, Erickson SE (2011) Distribution and abundance of introduced fishes in Florida's Charlotte Harbor estuary. *Gulf and Caribbean Research* 23: 13–22, <https://doi.org/10.18785/gcr.2301.02>
- Ishikawa T, Tachihara T (2014) Introduction history of non-native freshwater fish in Okinawa-jima Island: ornamental aquarium fish pose the greatest risk for future invasions. *Ichthyological Research* 6: 17–26, <https://doi.org/10.1007/s10228-013-0367-6>
- Jones RW, Weyl OLF, Swartz ER, Hill MP (2013) Using a unified invasion framework to characterize Africa's first loricariid catfish invasion. *Biological Invasions* 15: 2139–2145, <https://doi.org/10.1007/s10530-013-0438-7>
- Joshi RC (2006) Invasive alien species (IAS): concerns and status in the Philippines. In: *Proceedings of the International Workshop on the Development of Database (APASD) for Biological Invasion*, FFTC, Taichung, Taiwan, China, pp 1–23
- Jumawan JC, Herrera AA (2014) Ovary morphology and reproductive features of the female suckermouth sailfin catfish, *Pterygoplichthys disjunctivus* (Weber 1991) from Marikina River, Philippines. *Asian Fisheries Science* 27: 75–89
- Jumawan JC, Herrera AA (2015) Histological and ultrastructural characteristics of the testis of the invasive suckermouth sailfin

- catfish *Pterygoplichthys disjunctivus* (Siluriformes: Loricariidae) from Marikina River, Philippines. *Tissue and Cell* 47: 17–26, <https://doi.org/10.1016/j.tice.2014.10.005>
- Kaplan EL, Meier P (1958) Nonparametric estimation from incomplete observations. *Journal of the American Statistical Association* 53: 457–481, <https://doi.org/10.1080/01621459.1958.10501452>
- Kefford BJ, Papas PJ, Metzeling L, Nugegoda D (2004) Do laboratory salinity tolerances of freshwater animals correspond with their field salinity? *Environmental Pollution* 129: 355–362, <https://doi.org/10.1016/j.envpol.2003.12.005>
- Kestrup Å, Ricciardi A (2010) Influence of conductivity on the life history traits of exotic and native amphipods in the St. Lawrence River. *Fundamental and Applied Limnology* 176: 249–262, <https://doi.org/10.1127/1863-9135/2010/0176-0249>
- Keszka S, Panicz R, Tanski A (2008) First record of the Leopard Pleco, *Pterygoplichthys gibbiceps* (Actinopterygii, Loricariidae) in the Brda River in the centre of Bydgoszcz (northern Poland). *Acta Ichthyologica Et Piscatoria* 38: 135–138, <https://doi.org/10.3750/AIP2008.38.2.08>
- Knight JDM (2010) Invasive ornamental fish: a potential threat to aquatic biodiversity in peninsular India. *Journal of Threatened Taxa* 2: 700–704, <https://doi.org/10.11609/JoTT.o2179.700-4>
- Koohafkan P, Altieri MA (2011) Globally important agricultural heritage systems: legacy for the future. Food and Agriculture Organization of the United Nations, Rome, 41 pp
- Krishnakumar K, Raghavan R, Prasad G, Kumar AB, Sekharan M, Pereira B, Ali A (2009) When pets become pests – exotic aquarium fishes and biological invasions in Kerala, India. *Current Science* 97: 474–476
- Kumar AB (2000) Exotic fishes and freshwater fish diversity. *Zoos' Print Journal* 15: 363–367, <https://doi.org/10.11609/JoTT.ZPJ.15.11.363-7>
- Kumar AB, Smrithy R, Sureshkumar U, George S (2015) Invasion of South American Suckermouth armoured catfish *Pterygoplichthys* spp. (Loricariidae) in Kerala, India – a case study. *Journal of Threatened Taxa* 7: 6987–6995, <https://doi.org/10.11609/JoTT.o4133.6987-95>
- Kurup BM, Ranjeet K (2002) Integration of freshwater prawn culture with rice farming in Kuttanad, India. *Naga, World Fish Center Quarterly* 25: 16–19
- Liang SH, Wu HP, Shieh BS (2005) Size, structure, reproductive phenology and sex ratio of an exotic armoured catfish (*Liposacrus multiradiatus*) in Kaoping River of Southern Taiwan. *Zoological Studies* 44: 252–259
- Loftus WF (1988) Distribution and ecology of exotic fishes in Everglades National Park. In: Thomas LK (ed), Management of Exotic Species in Natural Communities. Proceedings of the George Wright Society Conference on Science in the National Parks, Vol 5. Ft. Collins, CO, pp 24–34
- Magalhães ALB, Jacobi CM (2013) Invasion risks posed by ornamental freshwater fish trade to southeastern Brazilian rivers. *Neotropical Ichthyology* 11: 433–441, <https://doi.org/10.1590/S1679-62252013005000003>
- Marr SM, Ellender BR, Woodford DJ, Alexander ME, Wasserman RJ, Ivey P, Zengeya T, Weyl OLF (2017) Evaluating invasion risk for freshwater fishes in South Africa. *Bothalia* 47: a 2177
- Martin GD, Vijay JG, Laluraj CM, Madhu NV, Joseph T, Nair M, Gupta GVM, Balachandran KK (2008) Fresh water influence on nutrient stoichiometry in a tropical estuary, southwest coast of India. *Applied Ecology and Environmental Research* 6: 57–64, https://doi.org/10.15666/aeer/0601_057064
- McIvor CC, Odum WE (1988) Food, predation risk, and microhabitat selection in a marsh fish assemblage. *Ecology* 69: 1341–1351, <https://doi.org/10.2307/1941632>
- Meena M, Sundaramanickam A, Ajith Kumar TT (2016) Occurrence of a *Pterygoplichthys disjunctivus* (Weber, 1991) population in Cauvery River System, Tamil Nadu, South India. *International Journal of Fisheries and Aquaculture* 8: 62–66, <https://doi.org/10.5897/IJFA2015.0526>
- Menon NN, Balchand AN, Menon NR (2000) Hydrobiology of the Cochin backwater system – a review. *Hydrobiologia* 430: 149–183, <https://doi.org/10.1023/A:1004033400255>
- Muralidharan M (2017) Do alien species matter? Impacts of invasions in Indian freshwater systems and challenges in management. *International Journal of Aquatic Biology* 5(2): 114–127
- Muralidharan M, Manikandan K, Gobi M (2015) Extended distribution of the invasive Sucker catfish *Pterygoplichthys pardalis* (Pisces: Loricariidae) to Cauvery river system of Peninsular India. *International Journal of Aquatic Biology* 3: 14–18
- Myers GS (1949) Usage of anadromous, catadromous and allied terms for migratory fishes. *Copeia* 1949: 89–97, <https://doi.org/10.2307/1438482>
- Nakabo T (2002) Fishes of Japan with Pictorial Keys to the Species, English edition, Tokai University Press, Tokyo, 1474 pp
- Nelson J (1994) Fishes of the World. John Wiley & Sons, Inc., New York, 600 pp
- Nico LG, Butt PL, Johnson GR, Jelks HL, Kail M, Walsh SJ (2012) Discovery of the South American suckermouth catfishes (Loricariidae, *Pterygoplichthys* spp.) in the Santa Fe River drainage, Suwannee River basin, USA. *Bioinvasions Records* 1: 179–200, <https://doi.org/10.3391/bir.2012.1.3.04>
- Nico LG, Loftus WF, Reid JP (2009) Interactions between non-native armoured suckermouth catfish (Loricariidae: *Pterygoplichthys*) and native Florida manatee (*Trichechus manatus latirostris*) in artesian springs. *Aquatic Invasions* 4: 511–519, <https://doi.org/10.3391/ai.2009.4.3.13>
- Nico LG, Martin TR (2001) The South American Suckermouth Armored Catfish, *Pterygoplichthys anisitsi* (Pisces: Loricariidae), in Texas, with comments on foreign fish introductions in the American Southwest. *Southwestern Naturalist* 46: 98–104, <https://doi.org/10.2307/3672381>
- Ozdilek SY (2007) Possible threat for middle-east inland water: an exotic and invasive species, *Pterygoplichthys disjunctivus* (Weber, 1991) in Asi river, Turkey. *Journal of Fisheries and Aquatic Sciences* 24(3–4): 303–306
- Paavola M, Olenin S, Leppäkoski E (2005) Are invasive species most successful in habitats of low native species richness across European brackish water seas? *Estuarine Coastal and Shelf Science* 64: 738–750, <https://doi.org/10.1016/j.ecss.2005.03.021>
- Page LM, Robins RH (2006) Identification of sailfin catfishes (Teleostei: Loricariidae) in south-eastern Asia. *The Raffles Bulletin of Zoology* 54: 455–457
- Panikkar P, Jagadeesh TD, Rao DSK, Sarkar UK, Naska M (2015) First record of non-native loricariid catfish, *Pterygoplichthys disjunctivus* (Weber, 1991) (Siluriformes, Loricariidae) in Cauvery River of peninsular India. *The Bioscan* 10: 1659–1663
- Parente TE, Moreira DA, Magalhães MGP, de Andrade PCC, Furtado C, Haas BJ, Stegeman JJ, Hahn ME (2017) The liver transcriptome of suckermouth armoured catfish (*Pterygoplichthys anisitsi*, Loricariidae): identification of expansions in defense gene families. *Marine Pollution Bulletin* 115: 352–361, <https://doi.org/10.1016/j.marpolbul.2016.12.012>
- Peterson MS, Meador MR (1994) Effects of salinity on freshwater fishes in coastal plain drainages in the Southeastern U.S. *Reviews in Fisheries Science* 2: 95–121, <https://doi.org/10.1080/10641269409388554>
- Piazzini S, Lori E, Favilli L, Cianfanelli S, Vanni S, Manganelli G (2010) Invasion note: a tropical fish community in thermal waters of southern Tuscany. *Biological Invasions* 12: 2959–2965, <https://doi.org/10.1007/s10530-010-9695-x>
- Raghavan R, Dahanukar N, Tlusty MF, Rhyne AL, Kumar KK, Molur S, Rosser AM (2013) Uncovering an obscure trade: threatened freshwater fishes and the aquarium pet markets. *Biological Conservation* 164: 158–169, <https://doi.org/10.1016/j.biocon.2013.04.019>
- Rao KR, Sunchu V (2017) A report on *Pterygoplichthys pardalis* Amazon sailfin sucker mouth catfishes in freshwater tanks at

- Telangana state, India. *International Journal of Fisheries and Aquatic Studies* 5: 249–254
- Rueda-Jasso RA, Campos-Mendoza A, Arreguín-Sánchez F, Díaz-Pardo E, Martínez-Palacios CA (2013) The biological and reproductive parameters of the invasive Armored Catfish *Pterygoplichthys disjunctivus* from Adolfo López Mateos El Infiernillo Reservoir, Michoacán-Guerrero, Mexico. *Revista Mexicana de Biodiversidad* 84: 318–326
- Sandilyan S (2016) Occurrence of ornamental fishes: a looming danger for inland fish diversity of India. *Current Science* 110: 2099–2104, <https://doi.org/10.18520/cs/v110/i11/2099-2104>
- Savage IR (1956) Contributions to the theory of rank order statistics – the two-sample case. *Annals of Mathematical Statistics* 27: 590–615, <https://doi.org/10.1214/aoms/1177728170>
- Schofield PJ, Nico LG (2009) Salinity tolerance of non-native Asian swamp eels (Teleostei: Synbranchidae) in Florida, USA: comparison of three populations and implications for dispersal. *Environmental Biology of Fishes* 85: 51–59, <https://doi.org/10.1007/s10641-009-9456-9>
- Scott DM, Rabineau J, Wilson RW, Hodgson DJ, Brown JA (2008) Can pikeperch colonise new freshwater systems via estuaries? Evidence from behavioural salinity tests. *Marine and Freshwater Research* 59: 694–702, <https://doi.org/10.1071/MF07241>
- Selvam V (2003) Environmental classification of mangrove wetlands of India. *Current Science* 84: 757–765
- Simonović P, Nikolić V, Grujić S (2010) Amazon Sailfin Catfish *Pterygoplichthys pardalis* (Castellnau, 1855) (Loricariidae, Siluriformes), a new fish species recorded in the Serbian section of the Danube River. Second Balkan Conference on Biology, 21-23 May 2010. Plovdiv, Biotechnology & Biotechnological Equipment 24/2010/ Special Edition/On-Line, pp 655–660
- Singh AK (2014) Emerging alien species in Indian aquaculture: prospects and threats. *Journal of Aquatic Biology & Fisheries* 2: 32–41
- Singh AK, Ansari A, Srivastava SC, Verma P, Pathak AK (2014) Impacts of invasive fishes on fishery dynamics of the Yamuna River, India. *Agricultural Sciences* 5: 813–821, <https://doi.org/10.4236/as.2014.510086>
- Singh AK, Kumar D, Srivastava SC, Ansari A, Jena JK, Sarkar UK (2013) Invasion and impacts of alien fish species in the Ganga River, India. *Aquatic Ecosystem Health & Management* 16: 408–414, <https://doi.org/10.1080/14634988.2013.857974>
- Singh AK, Lakra WS (2011) Risk and benefit assessment of alien fish species of the aquaculture and aquarium trade into India. *Reviews in Aquaculture* 3: 3–18, <https://doi.org/10.1111/j.1753-5131.2010.01039.x>
- Sinha RK, Sarkar UK, Lakra WS (2010) First record of the southern sailfin catfish, *Pterygoplichthys anisisti* Eigenmann & Kennedy, 1903 (Teleostei: Loricariidae), in India. *Journal of Applied Ichthyology* 26: 606–608, <https://doi.org/10.1111/j.1439-0426.2010.01474.x>
- Smith SA, Bell G, Bermingham E (2004) Cross-Cordillera exchange mediated by the Panama Canal increased the species richness of local freshwater fish assemblages. *Proceedings of the Royal Society London B: Biological Sciences* 271: 1889–1896, <https://doi.org/10.1098/rspb.2004.2796>
- Soundararajan N, Raj RM, Kamaladhasan N, Saidanyan RI, Chandrasekaran S (2015) On-line trade of aesthetic exotic organisms: sword of Damocles? *Current Science* 109: 1404–1410, <https://doi.org/10.18520/cs/v109/i8/1404-1410>
- Subramanian S (1988) Rice-fish culture: status and potential for increased production in the southwestern states of India. *Journal of the Indian Fisheries Association* 18: 103–108
- Sumanasinghe HPW, Amarasinghe US (2013) Population dynamics of accidentally introduced Amazon Sailfin Catfish, *Pterygoplichthys pardalis* (Siluriformes, Loricariidae) in Pologolla reservoir, Sri Lanka. *Sri Lanka Journal of Aquatic Sciences* 18: 37–45, <https://doi.org/10.4038/sljas.v18i0.7040>
- Villalba-Villalba AG, Ramírez-Suárez JC, Valenzuela-Soto EM, Sánchez GG, Ruiz GC, Pacheco-Aguilar R (2013) Trypsin from viscera of vermiculated sailfin catfish, *Pterygoplichthys disjunctivus*, Weber, 1991: its purification and characterization. *Food Chemistry* 141: 940–945, <https://doi.org/10.1016/j.foodchem.2013.03.078>
- Wei H, Copp GH, Vilizzi L, Liu F, Gu D, Luo D, Xu M, Mu X, Hu Y (2017) The distribution, establishment and life-history traits of non-native sailfin catfishes *Pterygoplichthys* spp. in the Guangdong Province of China. *Aquatic Invasions* 12: 241–249, <https://doi.org/10.3391/ai.2017.12.2.11>
- Welcomme R, Vidthayanom C (2003) The impacts of introductions and stocking of exotic species in the Mekong Basin and policies for their control. MRC Technical Paper No. 9, Mekong River Commission, Phnom Penh, 35 pp
- Wu LW, Liu CC, Lin SM (2011) Identification of exotic Sailfin Catfish species (*Pterygoplichthys*, Loricariidae) in Taiwan based on morphology and mtDNA sequences. *Zoological Studies* 50: 235–246
- Zworykin DD, Budaev SV (2013) Non-indigenous armoured catfish in Vietnam: invasion and systematic. *Ichthyological Research* 60: 327–333, <https://doi.org/10.1007/s10228-013-0356-9>

Supplementary material

The following supplementary material is available for this article:

Table S1. Records of *Pterygoplichthys* sp. from Kerala, India.

This material is available as part of online article from:

http://www.reabic.net/journals/mbi/2018/Supplements/MBI_2018_Kumar_etal_Table_S1.xlsx