

Research Article

Experimental evidence that the invasive snail *Potamopyrgus antipodarum* (Gray, 1843) survives passage through the digestive tract of common riverine fishRokas Butkus^{1,*} and Vytautas Rakauskas²¹Marine research institute, Klaipėda university, H. Manto st. 84, Klaipėda, Lithuania²Nature Research Centre, Akademijos st. 2, LT-08412 Vilnius, Lithuania

*Corresponding author

E-mail: butkus.rokas@gmail.com

Citation: Butkus R, Rakauskas V (2020) Experimental evidence that the invasive snail *Potamopyrgus antipodarum* (Gray, 1843) survives passage through the digestive tract of common riverine fish. *Management of Biological Invasions* 11(1): 96–104, <https://doi.org/10.3391/mbi.2020.11.1.07>

Received: 19 August 2019**Accepted:** 9 December 2019**Published:** 27 January 2020**Thematic editor:** Calum MacNeil**Copyright:** © Butkus and RakauskasThis is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International - CC BY 4.0](https://creativecommons.org/licenses/by/4.0/)).

OPEN ACCESS

Abstract

Potamopyrgus antipodarum is one of the most widespread invasive species worldwide and its expansion is ongoing. Although the snail has already been documented as surviving the digestive tract of various fish, there is insufficient data on potential riverine fish predators of *P. antipodarum* and of the ability of the snail to survive digestion by these fish. Lotic ecosystems are unfavourable for the upstream spread of this species by active movement and thus it is suspected that benthivorous riverine fish facilitate such upstream dispersion. In general, the results of this study indicated that *P. antipodarum* is not a preferable food object for the studied riverine fish. The study did indicate, however, that the invasive snail is capable of surviving the digestive tract of all the studied fish. As accidental consumption of abundant *P. antipodarum* with other food objects has already been documented, this study suggests that riverine fish can act as potential secondary dispersion vectors, facilitating the upstream dispersion of the snail.

Key words: zoochory, secondary dispersion vector, potential predators, gastrointestinal tract, upstream spread

Introduction

Throughout many parts of the world, lotic or running water ecosystems have been invaded by a variety of non-indigenous plant and animal taxa. Due to their connections with other aquatic habitats, such as lakes, streams and lagoons, these ecosystems are susceptible to non-native species. However, running waters are generally unfavourable for the active spread for most of these organisms, particularly in the upstream direction. Accelerating the dispersion of the various aquatic invaders, a variety of human- and non-human-mediated vectors have been documented in such ecosystems (Holmlund 2014; Green 2016; Coughlan et al. 2017 and others).

Potamopyrgus antipodarum (Gray, 1843) is a prosobranch snail native to New Zealand (Winterbourn 1970). In recent years, it has become one of the most widespread invasive aquatic species in the world and further rapid expansion is still ongoing (Radea et al. 2008; Son et al. 2008; Butkus et al. 2014 and others). The snail has been introduced and has established

in a wide range of aquatic ecosystems, including running waters (Ponder 1988; Bowler 1991; Alonso and Castro-Diez 2008; Davidson et al. 2008). Among the lotic habitats in which it has been recorded are streams flowing out from lakes, stream riffles, drainage ditches, high-elevation temperate streams, coastal waterways, Mediterranean-climate streams and cold water springs (Schreiber et al. 2002; Hall et al. 2003; Brown et al. 2008; Brenneis et al. 2011; Son et al. 2008; Butkus et al. 2014 and others). In these ecosystems, snail densities vary from scattered individuals in streams flowing out from lakes to extremely high densities of up to 500,000 ind. m⁻² observed in productive streams (Hall et al. 2006; Butkus *personal observation*). Dispersion in lotic waters is frequently mediated by both active and a number of passive dispersion vectors. Due to their small individual size (shell length up to 6–7 mm), the active spread of invading snails is slow in running waters. A study by Haynes et al. (1985) showed that *P. antipodarum* demonstrate positive rheotactic behaviour, while Zaranko et al. (1997) found that the snail can move upstream by up to 60 m over a period of three months. However, with current distribution patterns showing that the species is frequently recorded in a very short timespan at considerable distances from initial introduction points (Butkus et al. 2014, Butkus and Vaitonis 2019), it is likely that secondary dispersion vectors facilitate the spread of the invaders. According to previous studies, the transfer of *P. antipodarum* by human carriers, such as during water recreation, is one of the most significant agents in the dispersal of this invader in some areas (Richards et al. 2004; Holmlund 2014; Butkus and Vaitonis 2019). As previous studies indicated, *P. antipodarum* can survive desiccation for several days (Gaino et al. 2008) and thus the possibility of transport while attached to boats or on waterfowl feathers is high (Crozet et al. 1980). More importantly, several studies have also found that the invader can survive the digestive tract of marine and lake benthivorous fish or waterbirds (Aarnio and Bonsdorff 1997; Bruce and Moffitt 2009; Brenneis et al. 2011; Rakauskas et al. 2016). It has been shown that, to some degree, *P. antipodarum* is consumed by marine or brackish water benthivorous fish species in the field (Azour 2011; Brenneis et al. 2011; Rakauskas et al. 2013). In this way, fish may act as potential dispersion vectors and thus reduce colonization time in a particular ecosystem or even between different ecosystems. Most of the evaluated studies that analysed the ability of *P. antipodarum* to survive digestion focused mostly on fish common in still water ecosystems. By contrast, relatively few studies sought to identify riverine fish species that were capable of consuming the invader effectively (Cada 2004; Bersine et al. 2008; Brenneis et al. 2011). Furthermore, most of these studies focused on salmonid fish, these not being common or benthivorous in temperate running waters. Nevertheless, given that active dispersion of *P. antipodarum* is generally limited in lotic ecosystems, it can be assumed that riverine benthivorous fish species are potential dispersion vectors for the snail.

This paper investigates riverine fish species as a potential secondary dispersion vectors for *P. antipodarum* in lotic ecosystems. As some of the riverine benthivorous fish species are common and relatively abundant, there is a high probability that they act as secondary dispersion vectors and facilitate both downstream and upstream spread of *P. antipodarum*. However, due to a lack of data, the actual role of riverine fish in the secondary dispersion of the invasive snail is unclear. To plug this gap in current knowledge, we performed laboratory analysis of the consumption of *P. antipodarum* by the main local riverine benthivorous fish species. In the laboratory experiments, we tested whether these fish species consumed *P. antipodarum* in experimental aquariums in which the snails were the only available prey, and whether the snail survived passage through the digestive tract of these fish. In general, we predicted that *P. antipodarum* could survive passage through the digestive tract of most benthivorous riverine fish species, and thus these species could act as secondary dispersion vectors in river ecosystems.

Materials and methods

Individuals of *P. antipodarum* were collected in Lake Vilkokšnis using a standard dip net and sieving out the sand. The snails were transported to the laboratory and maintained in two 10 L plastic buckets at a temperature of 17 °C with part of the water changed every second day for a week before the experiments. Stones with epilythic algae collected in Lake Vilkokšnis were added as a food source for the snails. Snails with a shell length larger than 3 mm were selected and used in all experiments. Benthivorous fish species such as *Barbatula barbatula* (Linnaeus, 1758), *Barbus barbus* (Linnaeus, 1758), *Cottus gobio* Linnaeus, 1758, *Gobio gobio* (Linnaeus, 1758) and the long-distance migrant *Vimba vimba* (Linnaeus, 1758) were selected for the experiments. All of these are common benthivorous fish in running waters of central Europe, including Lithuania (Virbickas 2013, 2014). All fish specimens, except *V. vimba*, were collected in the River Neris (54°43'19.13"N; 25°02'56.33"E), using battery-powered electric fishing gear. Specimens of *V. vimba* were caught in the reservoir of Kaunas Hydroelectric Power Plant (54°53'54.15"N; 24°05'01.28"E) using multi-mesh benthic gillnets. It is worthy of note to mention that the invasive snail has not been recorded in either of the sampling sites. All fish specimens were acclimated in 92 L tanks filled with tap water circulating through ammonia filters for a week. During this period, frozen midge larvae were used as food. During the acclimation period and during the experiments, a natural photoperiod (18 h daylight, 6 h dark) and a temperature of 17 °C were maintained. The fish were not fed for 48 h prior to the experiments.

Two experiments were conducted in eight 9.5 L tanks connected to an ammonia filter forming a closed recirculating system – experiment 1 evaluated which of the riverine benthivorous fish consumed *P. antipodarum*, while

experiment 2 assessed snail survival on passing through the digestive tract of the fish. Following the methods described in Rakauskas et al. 2016, no substrate was added to the experimental tanks. In the feeding experiment, 100 living snails were presented for each tested fish specimen, allowing the fish to forage for 18 h. After this, they were removed and live snail specimens were carefully pumped out and counted, thereby allowing an estimate of the exact number of ingested snails. As the feeding experiments showed a low propensity among the study fish species to consume *P. antipodarum*, snails were inserted in earthworms *Lumbricus terrestris* (Linnaeus, 1758) for the survivorship experiments following Rakauskas et al. (2016). In total, 15 snails were served in 2–3 earthworms for each fish. As smaller fish specimens were unable to consume two large earthworms, the snails were served in three smaller earthworms. As the probability of accidental consumption of the snail together with preferable food objects by benthivorous fish is relatively high in the wild, this method of snail presentation in the current experiment imitates this process. After foraging for 18 h, the fish were placed individually into further experimental aquariums with a net (5 mm mesh) fastened about 3 cm above the bottom, this restricting the ability of the fish to reconsume the faecal pellets. Fish were left to defecate for 48 h at 17 °C and the faecal material from each aquarium was siphoned and sieved (50 µm mesh) every 8 h. All snails with intact shells were put into separate petri dishes filled with tap water for 12 h to recover from the digestion processes. The number and condition of the snails were assessed using a dissecting microscope. Assessment of condition was based on snail appearance and movement – dead snails either remained deep within their shells or, if their body was exposed outside of the shell, exhibited no movement when probed. Generally, surviving snails crawled around in the petri dishes.

Statistical analysis

As the data did not meet the normality assumption of parametric methods (Shapiro-Wilk's *W* tests, $P < 0.05$), non-parametric statistics were used. Kruskal-Wallis ANOVA tests were applied to test for species effect on the number of *P. antipodarum* ingested by different fish species and on the number of snails that survived passing through the gastrointestinal tracts of different predators. If necessary, multiple comparison of mean ranks was applied. Calculations were performed using Statistica 10.

Results

All studied fish species consumed the presented snails, though the ingestion rate varied among the species (Kruskal-Wallis ANOVA test: $H_{4, 45} = 28.92$, $P < 0.001$; Table 1). *Barbus barbuis* showed the greatest prevalence for feeding on the invasive snail: every *B. barbuis* individual used in the experiment

Table 1. Experimental ingestion of *P. antipodarum* by five main riverine benthivorous fish species: the number of investigated fish individuals of each particular species (n); the number of investigated fish individuals that consumed *P. antipodarum* served in earthworms (n_c); the total body length of fish (TL); the percentage of fish individuals that ingested *P. antipodarum* (P); the percentage of ingested *P. antipodarum* individuals (C); the number of ingested *P. antipodarum* individuals per fish (I). Values are mean \pm standard deviation.

Species	Consumption experiment					Survivorship experiment		
	n	TL (cm)	P (%)	C (%)	I (ind.)	TL (cm)	n	n_c
<i>B. barbatula</i>	15	10.1 \pm 0.9	13.3	0.2 \pm 0.6	1.5 \pm 0.7	11.3	5	1
<i>B. barbus</i>	8	23.6 \pm 1.0	100.0	10.3 \pm 6.1	10.3 \pm 6.1	23.6 \pm 1.1	5	5
<i>C. gobio</i>	5	6.5 \pm 2.2	80.0	3.0 \pm 2.5	3.8 \pm 2.2	7.2 \pm 1.4	5	2
<i>G. gobio</i>	8	10.2 \pm 0.6	37.5	0.8 \pm 1.2	2.0 \pm 1.0	10.2 \pm 0.9	5	2
<i>V. vimba</i>	10	21.0 \pm 1.1	80.0	2.2 \pm 1.9	2.8 \pm 1.7	21.0 \pm 1.2	7	7




Figure 1. The Number of ingested *P. antipodarum* individuals per fish (red bars) and the number of snails that survived passing through the digestive tract (green bars). Values are mean \pm standard deviation.

ingested them. However, on average, they ate only 10.3% of the snails available in the aquariums (Table 1). The applied multiple comparison on the number of consumed snails indicated significant ($P = 0.009$) differences only between *B. barbus* and *V. vimba* specimens. The prevalence of the other studied fish species for feeding on the invasive snail varied from 13.3% in *B. barbatula* to 80% in *C. gobio* and *V. vimba*. However, the ingestion rates by *B. barbatula*, *C. gobio*, *G. gobio* and *V. vimba* were low, these fish were eating less than 3% of the available snails and, on average, less than four snails per fish (Table 1).

Survival experiments indicated that some of the tested fish did not consume the earthworms with the inserted *P. antipodarum* (Figure 1). The experiments also yielded no significant differences in the number of *P. antipodarum* surviving after passing through the gastrointestinal tract of the different fish species (Kruskal-Wallis ANOVA test: $H_{4, 16} = 3.36$, $P = 0.49$). consumed snails were able to survive passing through the digestive tract of

all the studied fish species (Figure 1). the number of surviving snails ranged from 20% in *G. gobio* to 60% in *B. barbatula*. Additionally, none of the studied fish were able to crush the shell of *P. antipodarum*, thus resulting in the passage of all consumed snails undigested.

Discussion

Running water ecosystems do not favour the active spread of *P. antipodarum*. Secondary dispersion vectors, including fish, may facilitate species dispersion in such ecosystems. There is, however, limited current data on common riverine fish as potential vectors for the spread of this invasive species. In this study, Potential riverine fish predators of *P. antipodarum* were evaluated under laboratory experiments. Prior to our experiments, we assumed that the snail would be able to survive passing through the digestive tract of these fish and we also expected that *V. vimba*, *G. gobio* and *B. barbatus* could be potential predators of *P. antipodarum* and would consume a significant quantity of the presented snails. Molluscs are common food objects for these fish species in the wild (Kennedy and Fitzmaurice 1972; Okgerman et al. 2013; Djikanović et al. 2015) and, more importantly, gastropods (mainly of the genus *Hydrobia*) predominate in the diet of large *G. gobio* individuals in some ecosystems (Kennedy and Fitzmaurice 1972). Meanwhile, we expected *Cottus gobio* and *Barbatula barbatula* to consume insignificant quantities of the *P. antipodarum* individuals, as molluscs are not common food objects in the diets of these species (Hyslop 1982; Rău et al. 2015). Overall, the results of this study indicated that most of the studied riverine fish fed to a very limited degree on the *P. antipodarum* under laboratory experiments. This corresponds with previous results on benthivorous lake fish, whereby most of the studied fish consumed an insignificant number of the snails during laboratory experiments (Rakauskas et al. 2016). Despite several studies indicating benthivorous fish avoid consuming *P. antipodarum* (Cada 2004; Bersine et al. 2008; Rakauskas et al. 2016), the role of these species in facilitating snail dispersion can still be assumed important. the accidental consumption of the snail together with preferable food objects is highly possible and this probability to consume accidentally increases with increasing snail density. In the study of Hall et al. (2003), an extremely high density of up to 500,000 *P. antipodarum* ind. m⁻² were recorded in a river. In this case, the invasive snail dominated over other taxa and thus had a high probability to be consumed accidentally.

In this study, *P. antipodarum* was shown to survive digestion by riverine benthivorous fish under laboratory conditions, including by the migratory *V. vimba*. Such survival through the digestive tract, particularly in a migratory species, may lead to increased dispersion rates of the snail in river ecosystems. Currently, the translocation of aquatic invaders,

including *P. antipodarum*, both upstream and between river basins, has been widely attributed to anthropogenic activities (Alonso and Castro-Díez 2008; Kappes and Haase 2012; Banha et al. 2016). However, alternative natural vectors for passive dispersal remain under-researched (Kappes and Haase 2012; Banha et al. 2016; Coughlan et al. 2017). Zoochorous transport of one organism by another more mobile animal can facilitate dispersal of various taxa (Reynolds et al. 2015; Green 2016). Previous studies have indicated the ability of *P. antipodarum* to survive the gastrointestinal tract of various benthivorous fish under laboratory conditions (Aarnio and Bonsdorff 1997; Vinson and Baker 2008; Bruce and Moffitt 2009; Rakauskas et al. 2016). The results of this study correspond with these results. More importantly, previous studies have shown molluscs to be the predominant food source for the anadromous *V. vimba* (Bubinas and Vaitonis 2003; Ermolin and Shashulovskii 2006; Okgerman et al. 2013). In general, as *P. antipodarum* is capable of surviving the digestive tract of *V. vimba*, there is a high possibility of accelerated *P. antipodarum* dispersion. This has not, however, been directly documented yet.

Concluding remarks

The results of this study support the assumption that fish may have a potential role in the secondary dispersion of *P. antipodarum* in riverine ecosystems. However, this hypothesis needs additional investigation. Further research should be carried out to evaluate the proportion of the snail in riverine fish diet and the survival rates of the snail under natural conditions.

Acknowledgements

We thank Jos Stratford for revision of the English language in this paper and anonymous peer referees for valuable comments on an early draft of the article.

Funding declaration

This project has received funding from the European Social Fund (project No 09.3.3-LMT-K-712-02-0179) under grant agreement with the Research Council of Lithuania (LMTLT).

References

- Aarnio K, Bonsdorff E (1997) Passing the gut of juvenile flounder, *Platichthys flesus* (L.) - differential survival of zoobenthic prey species. *Marine Biology* 129: 11–14, <https://doi.org/10.1007/s002270050140>
- Alonso A, Castro-Díez P (2008) What explains the invading success of the aquatic mud snail *Potamopyrgus antipodarum* (Hydrobiidae, Mollusca)? *Hydrobiologia* 614: 107–116, <https://doi.org/10.1007/s10750-008-9529-3>
- Azour F (2011) Fødebiologi hos den sortmundede kutling *Neogobius melanostomus* i danske farvande. [Feeding ecology of the round goby *Neogobius melanostomus* in Danish waters]. BSc Thesis, University of Copenhagen [In Danish]
- Banha F, Gimeno I, Lanao M, Touya V, Durán C, Peribáñez M, Anastácio P (2016) The role of waterfowl and fishing gear on zebra mussel larvae dispersal. *Biological Invasions* 18: 115–125, <https://doi.org/10.1007/s10530-015-0995-z>
- Bersine K, Brenneis VEF, Draheim RC, Rub AMW, Zamon JE, Litton RK, Hinton SA, Sytsma MD, Cordell JR, Chapman JW (2008) Distribution of the invasive New Zealand mudsnail (*Potamopyrgus antipodarum*) in the Columbia River Estuary and its first recorded

- occurrence in the diet of juvenile Chinook salmon (*Oncorhynchus tshawytscha*). *Biological Invasions* 10: 1381–1388, <https://doi.org/10.1007/s10530-007-9213-y>
- Brown KM, Lang B, Perez KE (2008) The conservation ecology of North American pleurocerid and hydrobiid gastropods. *Journal of the North American Benthological Society* 27: 484–495, <https://doi.org/10.1899/07-062.1>
- Brenneis VEF, Sih A, de Rivera CE (2011) Integration of an invasive consumer into an estuarine food web: direct and indirect effects of the New Zealand mud snail. *Oecologia* 167: 169–179, <https://doi.org/10.1007/s00442-011-1962-8>
- Bruce RL, Moffitt CM (2009) Survival and passage of ingested New Zealand Mudsnailed through the intestinal tract of rainbow trout. *North American Journal of Aquaculture* 71: 287–301, <https://doi.org/10.1577/A08-033.1>
- Bubinas A, Vaitonis G (2003) The analysis of the structure, productivity and distribution of zoobenthocenoses in the Lithuanian economic zone of the Baltic Sea and the importance of some benthos species to fish diet. *Acta Zoologica Lituanica* 13: 114–124, <https://doi.org/10.1080/13921657.2003.10512555>
- Butkus R, Vaitonis G (2019) New records and distribution patterns of the invasive snail *Potamopyrgus antipodarum* (Gray, 1843) in Lithuanian inland waters. *BioInvasions Records* 8: 804–812, <https://doi.org/10.3391/bir.2019.8.4.08>
- Butkus R, Šidagyte E, Rakauskas V, Arbačiauskas K (2014) Distribution and current status of non-indigenous mollusc species in Lithuanian inland waters. *Aquatic Invasions* 9: 95–103, <https://doi.org/10.3391/ai.2014.9.1.08>
- Bowler P (1991) The rapid spread of the freshwater hydrobiid snail *Potamopyrgus antipodarum* (Gray) in the middle Snake river, southern Idaho. *Proceedings of Desert Fishes Council* 21: 173–182
- Cada C (2004) Interactions between the invasive New Zealand mudsnail, *Potamopyrgus antipodarum*, mayflies, and fish predators. MS thesis. Montana State University, Bozeman
- Coughlan NE, Stevens AL, Kelly TC, Dick JTA, Jansen MAK (2017) Zoochorous dispersal of freshwater bivalves: an overlooked vector in biological invasions? *Knowledge and Management of Aquatic Ecosystems* 418: 42, <https://doi.org/10.1051/kmae/2017037>
- Crozet B, Pedroli JC, Vaucher C (1980) Pre'mieres observations de *Potamopyrgus jenkinsi* (Smith) (Mollusca, Hydrobiidae) en Suisse romande. *Revue Suisse de Zoologie* 87: 807–811, <https://doi.org/10.5962/bhl.part.85552>
- Davidson T, Brenneis E, De Rivera C, Draheim R, Gillespie G (2008) Northern range expansion and coastal occurrences of the New Zealand mud snail, *Potamopyrgus antipodarum* (Gray, 1843) in the northeast Pacific. *Aquatic Invasions* 3: 349–353, <https://doi.org/10.3391/ai.2008.3.3.12>
- Djikanović V, Skorić S, Lenhardt M, Smederevac-Lalić M, Višnjić-Jeftić Z, Spasić S, Mićković B (2015) Review of sterlet (*Acipenser ruthenus* L. 1758) (Actinopterygii: Acipenseridae) feeding habitats in the River Danube, 1694–852 river km. *Journal of Natural History* 49: 411–417, <https://doi.org/10.1080/00222933.2013.877991>
- Ermolin VP, Shashulovskii VA (2006) On the results of *Vimba vimba* (Cypriniformes, Cyprinidae) introduction in the Volgograd Reservoir. *Journal of Ichthyology* 46: 487–489, <https://doi.org/10.1134/S0032945206060129>
- Gaino E, Scoccia F, Lancioni T, Ludovisi A (2008) The invader mudsnail *Potamopyrgus antipodarum* in the Tiber River basin (Central Italy). *Italian Journal of Zoology* 75: 253–261, <https://doi.org/10.1080/11250000701885513>
- Green AJ (2016) The importance of waterbirds as an overlooked pathway of invasion for alien species. *Diversity and Distribution* 22: 239–247, <https://doi.org/10.1111/ddi.12392>
- Haynes A, Taylor BR, Varley ME (1985) The influence of mobility of *Potamopyrgus jenkinsi* (Smith, E.A.) (Prosobranchia: Hydrobiidae) on its spread. *Archiv fur hydrobiologie* 103(4): 497–508
- Hall RO, Tank JL, Dybdahl MF (2003) Exotic snail dominate nitrogen and carbon cycling in a highly productive stream. *Frontiers in Ecology and Environment* 1: 407–411, [https://doi.org/10.1890/1540-9295\(2003\)001\[0407:ESDNAC\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2003)001[0407:ESDNAC]2.0.CO;2)
- Hall RO, Dybdahl MF, Vanderloop MC (2006) Extremely high secondary production of introduced snails in rivers. *Ecological Applications* 16: 1121–1131, [https://doi.org/10.1890/1051-0761\(2006\)016\[1121:EHSP01\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2006)016[1121:EHSP01]2.0.CO;2)
- Holmlund E (2014) Watershed Stewardship Program: Summary of Programs and Research. Adirondack Watershed Institute of Paul Smith's College. Watershed Stewardship Program Report # AWI 2014-01
- Hyslop EJ (1982) The feeding habits of 0+ stone loach, *Nemacheilus barbatulus* (L.), and bullhead, *Cottus gobio* L. *Journal of Fish Biology* 21: 187–196, <https://doi.org/10.1111/j.1095-8649.1982.tb03998.x>
- Kappes H, Haase P (2012) Slow, but steady: dispersal of freshwater molluscs. *Aquatic Sciences* 74: 1–14, <https://doi.org/10.1007/s00027-011-0187-6>
- Kennedy M, Fitzmaurice P (1972) Some aspects of the biology of gudgeon *Gobio gobio* (L.) in Irish waters. *Journal of Fish Biology* 4: 425–440, <https://doi.org/10.1111/j.1095-8649.1972.tb05690.x>

- Okgerman HC, Yardimci CH, Dorak Z, Yilmaz N (2013) Feeding ecology of vimba (*Vimba vimba* L., 1758) in terms of size groups and seasons in Lake Sapanca, northwestern Anatolia. *Turkish Journal of Zoology* 37: 288–297
- Ponder WF (1988) *Potamopyrgus antipodarum* - a Molluscan coloniser of Europe and Australia. *Journal of Molluscan Studies* 54: 271–285, <https://doi.org/10.1093/mollus/54.3.271>
- Radea C, Louvrou I, Economou-Amilli A (2008) First record of the New Zealand mud snail *Potamopyrgus antipodarum* J.E. Gray 1843 (Mollusca: Hydrobiidae) in Greece - notes on its population structure and associated microalgae. *Aquatic Invasions* 3: 341–344, <https://doi.org/10.3391/ai.2008.3.3.10>
- Rakauskas V, Pūtys Ž, Dainys J, Lesutiene J, Ložys L, Arbačiauskas K (2013) Increasing population of the invader round goby, *Neogobius melanostomus* (Actinopterygii: Perciformes: Gobiidae), and its trophic role in the Curonian Lagoon, SE Baltic Sea. *Acta Ichthyologica et Piscatoria* 43: 95–108, <https://doi.org/10.3750/AIP2013.43.2.02>
- Rakauskas V, Butkus R, Merkytė E (2016) Consumption of the invasive New Zealand mud snail (*Potamopyrgus antipodarum*) by benthivorous predators in temperate lakes: a case study from Lithuania. *Hydrobiologia* 775: 213–230, <https://doi.org/10.1007/s10750-016-2733-7>
- Rău MA, Plavan G, Strungaru S A, Nicoară M, Moglan I, Ureche D (2015) Feeding ecology of two sympatric fish species in a river ecosystem. *Analele Stiintifice Ale Universitatii Al I Cuza Din Iasi Sectiunea I Biologie Animala* Tom LXI: 11–18
- Reynolds C, Miranda NAF, Cumming GS (2015) The role of waterbirds in the dispersal of aquatic alien and invasive species. *Diversity and Distributions* 21: 744–754, <https://doi.org/10.1111/ddi.12334>
- Richards D, O’Connell P, Shinn DC (2004) Simple Control Method to Limit the Spread of the New Zealand Mudsnail *Potamopyrgus antipodarum*. *North American Journal of Fisheries Management* 24: 114–117, <https://doi.org/10.1577/M02-133>
- Schreiber ESG, Lake PS, Quinn GP (2002) Facilitation of native stream fauna by an invading species? Experimental investigation of the interaction of the snail, *Potamopyrgus antipodarum* (Hydrobiidae) with native benthic fauna. *Biological Invasions* 4: 317–325, <https://doi.org/10.1023/A:1020925022843>
- Son MO, Nabozhenko MV, Shokhin IV (2008) The Don River basin is a new stage of expansion of *Potamopyrgus jenkinsi* (Smith, 1889) (Gastropoda, Hydrobioidea) in Europe. *Doklady biological sciences* 419: 129, <https://doi.org/10.1134/S0012496608020178>
- Vinson MR, Baker MA (2008) Poor growth of rainbow trout fed New Zealand mudsnails *Potamopyrgus antipodarum*. *North American Journal of Fisheries Management* 28: 701–709, <https://doi.org/10.1577/M06-039.1>
- Virbickas T (2013) Ichtiofaunos tyrimai bei ekologines būklės pagal žuvų rodiklius įvertinimas Lietuvos upėse ir ežeruose [Fish fauna studies in Lithuanian rivers and lakes and their ecological status assessment based on the fish fauna metrics]. Report of Nature research centre, 76 pp [in Lithuanian]
- Virbickas T (2014) Žuvų tyrimai paviršiniuose telkiniuose ir jų ekologinės būklės įvertinimas pagal ichtiofaunos rodiklius [Fish fauna studies in a surface water bodies and their ecological status assessment based on the fish fauna metrics]. Report of Nature research centre, 97 pp [in Lithuanian]
- Winterbourn M (1970) The New Zealand species of the genus *Potamopyrgus* (Gastropoda: Hydrobiidae). *Malacologia* 10: 283–321
- Zaranko DT, Farara DG, Thompson FG (1997) Another exotic mollusc in the Laurentian Great Lakes: the New Zealand native *Potamopyrgus antipodarum* (Gray 1843) (Gastropoda, Hydrobiidae). *Canadian Journal of Fisheries and Aquatic Sciences* 54: 809–814, <https://doi.org/10.1139/f96-343>